

species is not commonly found in its range.

Moyar river originates in the Nilgiri hills at an altitude of about 1800 m and drains into the Lower Bhavani dam at 280 m. It cuts the Sigur plateau from the Mysore plateau to the north and forms a natural boundary between Tamil Nadu and Karnataka. It flows through the well-forested areas of Mudumalai Wildlife Sanctuary, Sigur reserve forest and Moyar Reserve Forest, for almost its full length. Hitherto, four individuals of *Nukta* species of fish have been recorded from this river. Occurrence of *Torkhudree*, a rare and threatened fish and *Puntius mudumalaiensis*, an endemic species of Moyar river of Mudumalai Wildlife Sanctuary here is remarkable. Downstream, poaching pressure is high and this river needs to be protected for its fish diversity.

Earlier records:

Sykes (1841), Day (1877, 1889), Hora (1942), Suter (1944), Kalawar and Kelkar (1956), Yazdani and Singh (1990), Singh (1990), Talwar and Jhingran

(1991) recorded this species from the rivers of Deccan, viz. River Inderanee, River Indrayani (type locality), River Krishna, Ujni wetland etc., but mostly from Maharashtra, and it has not been reported from Tamil Nadu earlier. Therefore, the present record of this fish in the Moyar river extends its range of distribution in southern India.

ACKNOWLEDGEMENT

Our sincere thanks to the Forest Department of Tamil Nadu for permission to work in the forests along Moyar river. Our thanks are also due to Mr. Kaliyappan, our efficient field assistant.

February 23, 1996

A. MANIMEKALAN
D.F. SINGH

*Sálim Ali Centre for Ornithology &
Natural History,
Kalampalayam P.O.,
Coimbatore-641 010.*

REFERENCES

- DAY, F. (1877): Fishes of India. Today & Tomorrow's Book Agency, New Delhi. 543 pp.
- _____. (1889): The Fauna of British India, including Ceylon and Burma. Fishes. Taylor and Francis, London. Vol. I: 548 pp.
- HORA, S.L. (1942): Notes on fishes in the Indian Museum. XLIII. On the systematic position of *Cyprinus nukta* Sykes. *Rec. Indian Mus.*, 44(1): 10-14.
- KALAWAR, A.G. & C.L. KELKAR (1956): Fishes of Kolhapur. *J. Bombay nat. Hist. Soc.* 53(4): 669-679.
- SINGH, D.F. (1990): Ichthyofauna of Maharashtra- Dhulia district. *Rec. Zool. Surv. India* 86(1): 83-91.
- SUTER, M. (1944): New record of fish from Poona. *J. Bombay nat. Hist. Soc.* 44(3): 408-414.
- SYKES, W.H. (1841): On the fishes of Dukhun. *Trans. Zool. Soc. Lond.* 2: 349, pl. LX-XLVIII.
- TALWAR, P.K. & A.G. JHINGRAN (1991): Inland Fishes. Vol. I: 297-298. Vol. I & II. Oxford & IBH Publishing Co. Pvt. Ltd., New Delhi, India.
- YAZDANI, G.M. & D.F. SINGH (1990): On the fish resources of Ujni wetland, Pune, Maharashtra, *J. Bombay nat. Hist. Soc.* 87: 157-160.

29. DUNG BEETLE (COLEOPTERA: SCARABAEIDAE: SCARABAEINAE) FAUNA OF BANGALORE, KARNATAKA

India, like other tropical countries, has a rich scarabaeid fauna, but in spite of overwhelming numbers they rarely make their presence felt. Not much work has been done on the fauna of this group of beetles after Arrow (1931) in his comprehensive account on Indian scarabs, reported four tribes, 26 genera and 354 species. A survey was conducted in and around Bangalore to study the dung beetle fauna

during 1981-1985. This survey revealed the presence of 61 species (Table 1) of Scarabaeinae belonging to three tribes. Of these, 33 are being reported for the first time from Bangalore. Members of the tribe Pannelini were not encountered. The great majority of the species belong to the genus *Onthophagus*. Seven species viz., *Heliocopris gigas*, *Onthophagus ramosellus*, *O. negligens*, *O. vividus*, *O. faveri*, *O.*

Table 1
DUNG BEETLE FAUNA AND THEIR ASSOCIATED MAMMALIAN EXCREMENT FROM BANGALORE, KARNATAKA, SOUTH INDIA.

No.	Species	Associated Mammal dung	No.	Species	Associated Mammal dung
Tribe Scarabaeini					
1.	<i>Scarabaeus (Kheper) sanctus</i> (Fabricius)	Sheep, Cow	31.	<i>C. volcanus</i> (Fabricius)	Cow, Sheep, Dog
2.	<i>S. gangeticus</i> (Castelnau)	Cow	* 32.	<i>C. unicornis</i> (Fabricius)	Cow
* 3.	<i>S. brahminius</i> Castelnau	Cow	* 33.	<i>C. inermis</i> Arrow	Cow, Dog
4.	<i>S. erichsoni</i> (Harold)	Cow, Sheep	34.	<i>C. indicus</i> Harold	Cow, Dog
* 5.	<i>Gymnopleurus cyaneus</i> (Fabricius)	Cow, Sheep	* 35.	<i>Onthophagus gazella</i> (Fabricius)	Cow, Elephant
6.	<i>G. spilotus</i> (Macleay)	Man, Cow, Sheep	36.	<i>O. recticornutus</i> Lansberge	Cow
7.	<i>G. koenigi</i> (Fabricius)	Sheep, Cow	* 37.	<i>O. duponti</i> Boucomont	Cow, Dog
* 8.	<i>G. dejeani</i> Castelnau	Cow	* 38.	<i>O. amplexus</i> Sharp	Cow
9.	<i>G. gemmatus</i> Harold	Man, Cow, Sheep	39.	<i>O. ramosus</i> (Wiedemann)	Cow
10.	<i>G. militaris</i> (Fabricius)	Man, Sheep, Cow	40.	<i>O. dama</i> (Fabricius)	Cow, Elephant, Horse
Tribe Sisyphini					
* 11.	<i>Sisyphus crispatus hirtus</i> (Wiedemann)	Sheep, Cow	41.	<i>O. pactolus</i> (Fabricius)	Cow, Sheep
* 12.	<i>S. longipes</i> (Oliver)	Sheep, Cow	* 42.	<i>O. unifasciatus</i> Schaller	Cow, Sheep, Dog, and carcass of Cow, Frog, Tenebrionid Beetle
* 13.	<i>S. neglectus</i> Gory	Sheep, Cow, Monkey			
14.	<i>S. hirtus</i> Wiedemann	Sheep, Cow	* 43.	<i>O. turbatus</i> Walker	Cow, Horse
Tribe Coprini					
* 15.	<i>Helicopriss bucephalus</i> (Fabricius)	Elephant, Cow	* 44.	<i>O. spinifex</i> (Fabricius)	Cow
16.	<i>H. gigas</i> (Linnaeus)	Elephant, Cow	45.	<i>O. quadridentatus</i> (Fabricius)	Cow, Elephant
17.	<i>Copris signatus</i> Walker	Sheep	46.	<i>O. igneus</i> Vigors	Elephant
18.	<i>C. repertus</i> Walker	Cow, Elephant	47.	<i>O. pygmaeus</i> (Schaller)	Dog, Cow, Sheep
* 19.	<i>C. fricator</i> Fabricius	Cow, Sheep	* 48.	<i>O. tarandus</i> (Fabricius)	Cow
20.	<i>C. andrewesi</i> Waterhouse	Sheep	* 49.	<i>O. centricornis</i> (Fabricius)	Horse, Cow
21.	<i>C. indicus</i> Gillet	Cow	* 50.	<i>O. laevigatus</i> (Fabricius)	Cow
22.	<i>Catharsius molossus</i> (L.)	Cow, Elephant, Pig	* 51.	<i>O. ludio</i> Boucomont	Cow, Sheep
23.	<i>C. pithecius</i> (Fabricius)	Cow	* 52.	<i>O. pusillus</i> (Fabricius)	Cow, Sheep
24.	<i>Onitis philemon</i> F.	Cow, Elephant	* 53.	<i>O. tritinctus</i> Boucomont	Sheep, Cow, Dog
25.	<i>O. subopacus</i> Arrow	Cow	54.	<i>O. ephippioderus</i> Arrow	Cow
* 26.	<i>O. siva</i> Gill	Elephant	55.	<i>O. kchatriya</i> Boucomont	Cow, carcass of Cow, Sheep
* 27.	<i>Drepanocerus setosus</i> (Wiedemann)	Cow, Elephant	* 56.	<i>O. gratus</i> Arrow	Cow, Sheep
* 28.	<i>Oniticellus pallipes</i> (Fabricius)	Cow	* 57.	<i>O. abreui</i> Arrow	Cow
* 29.	<i>O. cinctus</i> (Fabricius)	Cow	58.	<i>Onthaphagus</i> sp.	Cow
30.	<i>Caccobius meridionalis</i> Boucomont	Cow, Sheep, Dog, Pig	* 59.	<i>Phalops divisus</i> (Wiedemann)	Cow
			* 60.	<i>Liatongus rhadamistus</i> (Fabricius)	Cow
			* 61.	<i>Tiniocellus modestus</i> Arrow	Cow, Elephant

Note: * indicates species being reported from Bangalore for the first time.

brevicollis and *O. brahma* reported by Arrow (1931) from Bangalore were not found during this study.

ACKNOWLEDGEMENT

We thank Dr. R. Madge of the Natural History Museum (London) for helping us to identify the dung beetles.

February 23, 1996

K. VEENAKUMARI
Central Agricultural Research Institute,
P.B. No. 181, Port Blair, Andamans.

G. K. VEERESH
Vice Chancellor,
University of Agricultural Sciences,
G.K.V.K., Bangalore, 560065.

REFERENCES

ARROW, G.J., (1931): The Fauna of British India. Coleoptera, Lamellicornia, Part III, (Coprinae). Today and Tomorrow's Printers and Publishers, New Delhi.

30. FIRST RECORD OF *DIRHINUS ALTICORNIS* (MASI) AND *ANNECKEIDA ANGUSTIFRONS* BOUCEK (HYMENOPTERA: CHALCIDOIDEA) FROM INDIA

(With three text-figures)

During the faunal exploration of tropical rainforests of Western Ghats by Zoological Survey of India, Western Ghats Field Research Station, Calicut, two interesting chalcids were collected from semievergreen forest patches in Coorg district (Karnataka) and Kannur district (Kerala).

Dirhinus Dalman, one of the most distinctive genera of the family Chalcididae is distributed in all warmer countries of the world, Africa, Europe, Asia, Australia and Pacific islands. Members of this genus are parasitic on puparia of various Diptera, especially Calliphoridae, Sarcophagidae, Muscidae and also of certain Tephritidae.

Dirhinus alticornis (Masi), a remarkable species of the genus was originally described from Philippines by Masi (1927) under the name *Parenia alticornis*. Narendran (1989) examined a male specimen of *alticornis* from Philippines. One female specimen of the species was collected by me from a semievergreen forest patch at Aniyad, falling under the Kannavam RF of Kannur district, Kerala. The present record of *D. alticornis* (Masi) from the Western Ghats proves the further extension of its distribution to peninsular India and the third record from the Oriental Region.

D. alticornis is a characteristic species having the anterior inner edge of its frontal horn crenulate with an additional tooth on the outer edge. It has a strong facial tooth, and the posterior median area of

pronotum depressed with an impunctate shagreened area. In males the antenna is peculiar with a spatulate club.

Specimen examined: 1 FEMALE. INDIA: Kerala, Aniyad (Kannavam RF), 1. ii. 1995, Coll. P.M. Sureshan *et al.*

Anneckeida Boucek, an African genus of Torymidae is represented by four Oriental species from West Malaysia, Thailand, Laos, and East Malaysia, apart from the type species from Rhodesia (Africa). *A. angustifrons* was originally described by Boucek (1978) based on a female specimen collected from Thailand. One female specimen of this species was collected by me from a forest patch at Chitekanum, falling under the Sampage reserve forests of Coorg district, Karnataka. This record constitutes its rediscovery from the Oriental Region subsequent to the original description and proves the extension of its distribution to Peninsular India.

Like all other Oriental species *A. angustifrons* also has hind femur with a ventral comb of teeth which begins with a conspicuous larger tooth. The species is also characterised by a face with inner orbits, distinctly converging upwards frons only 0.25 x the breadth of head and the ocelli in acute angular triangle, with lateral ones virtually touching the eyes.

Specimen examined: 1 FEMALE. INDIA: Karnataka, Chitekanum (Samapge R.F.), 4.iii. 1994, Coll. P.M. Sureshan *et al.*